

YOUR GUIDE TO TREATMENT

STEP BY STEP

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.


LEMTRADA[®]
alemtuzumab ^{12mg} iv

BEFORE TREATMENT

DURING TREATMENT

AFTER TREATMENT

FAQS AND IMPORTANT
SAFETY INFORMATION

PLANNING FOR WHAT'S NEXT

Reading the information in this guide will help with your preparation before, during, and after your LEMTRADA treatment.

INDICATION

LEMTRADA is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include relapsing-remitting disease and active secondary progressive disease, in adults. Since treatment with LEMTRADA can increase your risk of getting certain conditions and diseases, LEMTRADA is generally prescribed for people who have tried 2 or more MS medicines that have not worked well enough. LEMTRADA is not recommended for use in patients with clinically isolated syndrome (CIS). It is not known if LEMTRADA is safe and effective for use in children under 17 years of age.

SELECTED IMPORTANT SAFETY INFORMATION

LEMTRADA can cause serious side effects including autoimmune problems, infusion reactions, stroke, tears in your arteries that supply blood to your brain (carotid and vertebral arteries), some kinds of cancers, thyroid problems, low blood counts (cytopenias), inflammation of the liver, hemophagocytic lymphohistiocytosis, Adult Onset Still's Disease, thrombotic thrombocytopenic purpura, autoimmune encephalitis, bleeding disorder (acquired hemophilia A), serious infections, progressive multifocal leukoencephalopathy (PML), inflammation of the gallbladder without gallstones (acalculous cholecystitis), and swelling of lung tissue (pneumonitis). Because of the risks of autoimmune problems, infusion reactions and some kinds of cancers, LEMTRADA is only available through a restricted program called the Risk Evaluation and Mitigation Strategy (REMS) Program.

WHAT'S DONE



DETERMINE WITH HCP IF LEMTRADA IS RIGHT FOR ME



ENROLL IN LEMTRADA REMS



READY TO BEGIN TAKING LEMTRADA

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

LEMTRADA[®]
alemtuzumab^{12mg} iv

EVERY STEP OF THE WAY THERE'S SUPPORT

It's comforting to have someone to talk to. Your *MS One to One*® Case Manager will be available, after you sign the LEMTRADA Services Form, to help you with:

- ✓ Verifying insurance benefits and answering other questions
- ✓ Locating an infusion center
- ✓ Organizing and preparing for your treatment
- ✓ Explaining what to expect after your infusion days
- ✓ Setting reminders for your monthly monitoring

FINANCIAL SUPPORT

To help make treatment costs and co-pays more affordable, the **LEMTRADA Co-Pay Program offers financial assistance to help eligible,* commercially insured patients receive:**

\$0 OUT-OF-POCKET COST
FOR LEMTRADA

\$100 UP TO PER DAY FOR CHARGES
RELATED TO THE
ADMINISTRATION OF
YOUR LEMTRADA INFUSION

*Patients who have coverage or prescriptions paid for in part or in full under Medicare, Medicaid, or other state or federally funded healthcare programs are not eligible. The LEMTRADA Co-Pay Program is subject to termination or modification at any time. If you are not eligible for the LEMTRADA Co-Pay Program and need help with out-of-pocket expenses, *MS One to One* can help review your coverage options. Depending on your specific situation, your *MS One to One* Case Manager can direct you to other patient assistance programs that may offer you financial support. Treatment-related, infusion out-of-pocket costs are not reimbursable in MA, MI, MN, or RI.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

MS ONE TO ONE® CASE MANAGERS ARE AVAILABLE 24/7

MS One to One will provide you with access to a case manager anytime. Regular call center hours are Mon-Fri, 8:30 am-8:00 pm ET, at **1-855-676-6326**. Any calls made after regular call center hours will be returned within 30 minutes.



name

phone number



name

address

phone number



Even if your dedicated *MS One to One* Case Manager is not available, someone is always there to answer questions.

LEMTRADA
alemtuzumab^{12mg} iv

JOURNEY TO TREATMENT CHECKLIST

6 WEEKS BEFORE



SCHEDULE APPOINTMENTS

Schedule your LEMTRADA infusion appointments. Ask about policies on bringing food, entertainment, visitors, etc.



CALL YOUR CASE MANAGER

Call *MS One to One*® at 1-855-676-6326 and give your case manager your appointment dates



GET PRETREATMENT TESTS AND VACCINATIONS

Complete blood and urine tests and a skin check as prescribed. Get any other tests and vaccinations your healthcare team has recommended



MAKE PLANS

Arrange for transportation, work coverage, and/or child care for days of infusion

1 WEEK BEFORE



CONFIRM DATES

Confirm your infusion appointments with your infusion center



GET MEDICATIONS

Get medications as prescribed by your healthcare team



CONFIRM PLANS

Confirm transportation to and from your infusion appointments. Do not drive yourself

1 DAY BEFORE



PACK

- headphones
- book
- food/snacks
- bottled water or refillable container



DOWNLOAD

- movies
- podcasts
- music playlist

MEDICATIONS YOUR DOCTOR WILL PRESCRIBE INCLUDE:

Corticosteroids to lower your chances of getting serious infusion reactions, your healthcare provider will give you a medicine called corticosteroids before your first 3 infusions of a treatment course. **Antihistamines and fever reducers** may be given prior to your infusion to reduce the chances of infusion reactions, or to treat them if they happen. **Antivirals** will be prescribed to you to take starting on your first day of infusions. Continue taking medications as recommended by your healthcare team.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.



LEARN HOW OTHER LEMTRADA PATIENTS PREPARED FOR THEIR TREATMENT JOURNEY

LEMTRADA®
alemtuzumab^{12mg} iv

TODAY IS TREATMENT DAY

You've done your research, spoken up to advocate for your treatment goals, and made all necessary arrangements.

Relax and remember...



TREATMENT CHECKLIST

✓
PACK BAG

✓
WEAR COMFY
CLOTHES

✓
TAKE RECOMMENDED
PRE-MEDICATIONS

✓
ARRIVE EARLY

✓
STAY HYDRATED

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

LEMTRADA[®]
alemtuzumab^{12mg} iv

STEPS TO HELP MANAGE INFUSION REACTIONS

LEMTRADA can cause serious infusion reactions that may cause death. Serious infusion reactions may happen while you receive, or up to 24 hours or longer after you receive LEMTRADA.

- **92%** of patients experienced infusion reactions in the LEMTRADA clinical trials
- **3%** of these reactions were serious



MANAGE INFUSION REACTIONS

You will receive your infusion at a healthcare facility with equipment and staff trained to manage infusion reactions, including serious allergic reactions and urgent heart or breathing problems.



MONITORING INFUSION REACTIONS

You will be watched while you receive, and for 2 hours or longer after you receive, LEMTRADA on each infusion day. If a serious infusion reaction happens while you are receiving LEMTRADA, your infusion may be stopped.



REMINDER

Make sure you stay hydrated before, during, and after your infusion.



**HEAR WHAT THE LEMTRADA
INFUSION PROCESS IS LIKE
FROM REAL-LIFE PATIENTS**

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

WATCH FOR THESE SYMPTOMS

Tell your healthcare team right away if you have any of the following symptoms of a serious infusion reaction during or after treatment:



SWELLING IN YOUR MOUTH OR THROAT



TROUBLE BREATHING



WEAKNESS



FAST, SLOW, OR IRREGULAR HEARTBEAT



CHEST PAIN



RASH

Some people have had serious and sometimes deadly strokes and tears in their carotid or vertebral arteries within 3 days of receiving LEMTRADA. Get help right away if you have any of the following symptoms that may be signs of a stroke or tears in your carotid or vertebral arteries: drooping of parts of your face, weakness on one side, sudden severe headache, difficulty with speech, or neck pain.

LEMTRADA[®]
alemtuzumab^{12mg}

YOUR TREATMENT COURSE IS **COMPLETE!**

Self-care is top priority in the days and months following your infusion.

Take the time to...

POST-TREATMENT TO-DO

- ✓ **REST, RELAX,
AND TAKE CARE
OF YOURSELF**
- ✓ **SCHEDULE
MONITORING
APPOINTMENTS**
- ✓ **PERFORM SYMPTOM
SELF-CHECKS**

**TAKE
CARE OF
YOU**



Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

LEMTRADA[®]
alemtuzumab^{12mg} IV

CARE FOR YOURSELF AFTER YOUR INFUSIONS

Be sure to follow any post-treatment instructions from your healthcare team, which may include:

- ✓ Washing hands frequently to reduce risk of infections
- ✓ Staying hydrated
- ✓ Resting
- ✓ Eating regularly
- ✓ Avoiding or adequately heating foods that may contain *Listeria* during and after LEMTRADA treatment*

SELECTED IMPORTANT SAFETY INFORMATION

LEMTRADA can cause serious side effects including autoimmune problems, infusion reactions, stroke, tears in your arteries that supply blood to your brain (carotid and vertebral arteries), some kinds of cancers, thyroid problems, low blood counts (cytopenias), inflammation of the liver, hemophagocytic lymphohistiocytosis, Adult Onset Still's Disease, thrombotic thrombocytopenic purpura, autoimmune encephalitis, bleeding disorder (acquired hemophilia A), serious infections, progressive multifocal leukoencephalopathy (PML), inflammation of the gallbladder without gallstones (acalculous cholecystitis), and swelling of lung tissue (pneumonitis). Because of the risks of autoimmune problems, infusion reactions and some kinds of cancers, LEMTRADA is only available through a restricted program called the Risk Evaluation and Mitigation Strategy (REMS) Program.

**Listeria* infections have occurred as early as 3 days after treatment and up to 8 months after the last LEMTRADA dose. For more information, please see page 10.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

TAKE AN ACTIVE ROLE IN COMPLETING YOUR MONITORING

Certain side effects of LEMTRADA can happen months and even years after treatment. That's why it is important to begin monitoring after your first infusion and continue regular monitoring until 4 years after your last infusion. Monitoring can help your healthcare team find potential side effects early.



Monthly blood and urine tests



Ongoing symptom self-checks



A yearly skin examination

MANY OPTIONS FOR MONTHLY MONITORING

MS One to One[®] offers a range of options for monthly blood and urine testing at no cost to you.



LAB: Any local LabCorp[®] or Quest Diagnostics[®] patient service center[†]



HEALTHCARE PROVIDER'S OFFICE: Samples can be taken at your healthcare team's office. With this option, you may incur a co-pay for your doctor's office visit. However, the cost of analyzing the samples will be covered by Sanofi.



LISTEN TO REAL-LIFE PATIENTS
TALK ABOUT THEIR EXPERIENCES
WITH LEMTRADA SUPPORT SERVICES

[†]Trademarks not owned by Sanofi corporation are the property of their respective owners.

LEMTRADA[®]
alemtuzumab^{12mg}iv

IDENTIFY SYMPTOMS

WITH SELF-CHECKS

Spot any physical changes by performing regular body checks at home by yourself in a mirror or with the help of a family member or friend.

EYES, NOSE, AND MOUTH

- Yellowing of the whites of your eyes
- Eye issues, including swelling and/or vision problems
- Bleeding from gums or nose that is new or takes longer than usual to stop
- Swelling in your mouth or throat
- Drooping of parts of your face

PAIN

- Neck pain
- Chest pain or tightness
- Sudden severe headache
- Stomach pain or discomfort

BOWEL AND URINARY ISSUES

- Constipation
- Blood in the urine (red or tea-colored)
- Decrease in urine
- Dark urine

SKIN


- Easy bruising
- Small, scattered spots that are red, pink, or purple
- Yellowing of the skin
- Bleeding from a cut that is hard to stop
- Swelling in your legs or feet
- Swelling in your neck
- New lump
- Rash

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.



REMINDER

Contact your healthcare team if you experience any of these symptoms, or new or worsening medical problems that have lasted several days including changes in energy or mood, problems with thinking and/or balance.



RESPIRATORY AND THROAT

- Cough that is not caused by a cold
- Coughing up blood
- Trouble swallowing or breathing
- Hoarseness or other voice changes that don't go away
- Swelling in your throat
- Wheezing
- Shortness of breath

OTHER

- Fast, slow, or irregular heartbeat
- Excessive sweating or feeling cold
- Weakness
- Problems using arms or legs
- Worsening tiredness
- Nervousness
- Unexplained weight loss or gain
- Loss of appetite
- Fever
- Swollen glands
- For women: heavier menstrual periods than normal
- Nausea or vomiting
- Weakness on one side
- Difficulty with speech
- Increased fatigue

YOU HAVE QUESTIONS WE HAVE ANSWERS

HOW LONG DOES EACH INFUSION LAST?

Each infusion takes approximately 4 hours, and your healthcare team will monitor you for at least 2 hours for infusion reactions.

WHAT KINDS OF SERIOUS SIDE EFFECTS CAN HAPPEN DURING AN INFUSION?

Tell your healthcare provider right away if you have any of the following symptoms of a serious infusion reaction during the infusion and after you have left the healthcare facility: swelling in your mouth or throat, trouble breathing, weakness, fast, slow or irregular heartbeat, chest pain or rash.

IS THERE ANYTHING I CAN DO TO HELP ADDRESS INFUSION SIDE EFFECTS?

- Your healthcare provider will give you a medicine called a corticosteroid before your first 3 infusions of a treatment course
- You may also be given other medicines before or after the infusion to try to reduce your chances of having these reactions or to treat them if they happen. If something feels out of the ordinary, contact your healthcare provider right away

WHY SHOULD I BE CONCERNED ABOUT POTENTIAL FOOD SOURCES OF LISTERIA?

People who receive LEMTRADA have an increased chance of getting an infection caused by the bacteria *Listeria*, which can lead to significant complications or death. Avoid foods that may be a source for *Listeria* (for example, deli meat, unpasteurized milk and cheese products, soft cheeses, or undercooked meat, seafood, or poultry), or make sure that the food you eat that may contain *Listeria* is heated well if you receive treatment with LEMTRADA.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.



Be confident that you are ready to take this next step in your treatment journey.

WHAT SHOULD I BRING TO MY INFUSION CENTER?

It is recommended that you bring a sweatshirt or a blanket and wear comfortable clothes. You should pack something to entertain yourself during your treatment.

- Additional tips to help you pass the time during treatment can be found on page 4
- Be sure to call the infusion center to be certain what food, drinks, and entertainment are permitted and ask questions like whether water will be available for refillable bottles

IS IT OKAY IF I GET PREGNANT WHILE TAKING LEMTRADA?

Before receiving LEMTRADA, tell your healthcare team if you are pregnant or plan to become pregnant, as it can harm your unborn baby.

- Use effective birth control to avoid pregnancy during a treatment round
- Continue to use birth control for at least 4 months after a treatment round

WHERE CAN I FIND ADDITIONAL SUPPORT?

An *MS One to One*® Case Manager is available 24/7* to provide support if you're feeling nervous or have questions before, during, and after treatment.

AFTER MY FIRST 2 ROUNDS, WHEN MIGHT I REQUIRE ADDITIONAL INFUSIONS?

After your initial 2 rounds, additional infusions are available, if needed, at least 1 year after your last treatment. Of the 387 patients who were in the Study 1 extension, 50% did not receive additional rounds of LEMTRADA or any other disease-modifying therapy (DMT) through 6 years. 29% received 1 additional round, 13% received 2 additional rounds, 2% received 3 additional rounds, 1% received 4 additional rounds, and 5% received another DMT and no additional rounds of LEMTRADA.†

*As a member of *MS One to One*, you'll have access to an on-duty case manager 24/7. Regular *MS One to One* call center hours are Mon-Fri, 8:30 am-8:00 pm ET.

†Information beyond 3 treatment courses is limited.



IMPORTANT SAFETY INFORMATION

LEMTRADA can cause serious side effects including:

Serious autoimmune problems: Some people receiving LEMTRADA develop a condition where the immune cells in your body attack other cells or organs in the body (autoimmunity), which can be serious and may cause death. Serious autoimmune problems may include:

- Immune thrombocytopenic purpura (ITP), a condition of reduced platelet counts in your blood that can cause severe bleeding that may cause life-threatening problems. Call your healthcare provider (HCP) right away if you have any of the following symptoms: easy bruising, bleeding from a cut that is hard to stop, coughing up blood, heavier menstrual periods than normal, bleeding from your gums or nose that is new or takes longer than usual to stop, small, scattered spots on your skin that are red, pink, or purple.
- Kidney problems called anti-glomerular basement membrane disease, which, if not treated, can lead to severe kidney damage, kidney failure that needs dialysis, a kidney transplant, or death. Call your HCP right away if you have any of the following symptoms: swelling of your legs or feet, blood in the urine (red or tea-colored urine), decrease in urine, fatigue, coughing up blood.

It is important for you to have blood and urine tests before you receive, while you are receiving and every month for 4 years or longer, after you receive your last LEMTRADA infusion.

Serious infusion reactions: LEMTRADA can cause serious infusion reactions that may cause death. Serious infusion reactions may happen while you receive, or up to 24 hours or longer after you receive LEMTRADA.

- You will receive your infusion at a healthcare facility with equipment and staff trained to manage infusion reactions, including serious allergic reactions, and urgent heart or breathing problems. You will be watched while you receive, and for 2 hours or longer after you receive, LEMTRADA. If a serious infusion reaction happens while you are receiving LEMTRADA, your infusion may be stopped.

Tell your HCP right away if you have any of the following symptoms of a serious infusion reaction during the infusion, and after you have left the healthcare facility:

- swelling in your mouth or throat
- trouble breathing
- weakness
- fast, slow, or irregular heartbeat
- chest pain
- rash

To lower your chances of getting a serious infusion reaction, your HCP will give you a medicine called corticosteroids before your first 3 infusions of a treatment course. You may also be given other medicines before or after the infusion to try to reduce your chances of having these reactions or to treat them if they happen.

Stroke and tears in your arteries that supply blood to your brain (carotid and vertebral arteries): Some people have had serious and sometimes deadly strokes and tears in their carotid or vertebral arteries within 3 days of receiving LEMTRADA. Get help right away if you have any of the following symptoms that may be signs of a stroke or tears in your carotid or vertebral arteries: drooping of parts of your face, weakness on one side, sudden severe headache, difficulty with speech, neck pain.

Certain cancers: Receiving LEMTRADA may increase your chance of getting some kinds of cancers, including thyroid cancer, skin cancer (melanoma), and blood cancers called lymphoproliferative disorders and lymphoma. Call your HCP if you have the following symptoms that may be a sign of thyroid cancer: new lump, swelling in your neck, pain in front of neck, trouble swallowing or breathing, hoarseness or other voice changes that do not go away, cough that is not caused by a cold.

Have your skin checked before you start receiving LEMTRADA and each year while you are receiving treatment to monitor for symptoms of skin cancer.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

LEMTRADA[®]
alemtuzumab^{12mg} iv

IMPORTANT SAFETY INFORMATION (CONTINUED)

Because of risks of autoimmunity, infusion reactions, and some kinds of cancers, LEMTRADA is only available through a restricted program called the LEMTRADA Risk Evaluation and Mitigation Strategy (REMS) Program.

Do not receive LEMTRADA if you:

- are allergic to alemtuzumab or to any of the inactive ingredients in LEMTRADA
- are infected with human immunodeficiency virus (HIV)
- have an active infection

Thyroid problems: Some patients taking LEMTRADA may get an overactive thyroid (hyperthyroidism) or an underactive thyroid (hypothyroidism). Call your HCP if you have: excessive sweating, unexplained weight loss, unexplained weight gain, fast heartbeat, eye swelling, nervousness, feeling cold, worsening tiredness, constipation.

Low blood counts (cytopenias): LEMTRADA may cause a decrease in some types of blood cells. Some people with these low blood counts have increased infections. Call your doctor right away if you have symptoms of cytopenias such as: weakness, chest pain, yellowing of the skin or whites of the eyes (jaundice), dark urine, fast heartbeat.

Inflammation of the liver: Call your HCP right away if you have symptoms such as unexplained nausea, stomach pain, tiredness, loss of appetite, yellowing of skin or whites of eyes, or bleeding or bruising more easily than normal.

Hemophagocytic lymphohistiocytosis: LEMTRADA may increase the risk of overactivity of the immune system that can be fatal if not diagnosed and treated early. If you experience symptoms such as fever, swollen glands, or skin rash, contact your HCP right away.

Adult Onset Still's Disease (AOSD): LEMTRADA may cause AOSD, a rare condition that can cause a high fever lasting more than 1 week, pain, stiffness with or without swelling in multiple joints, and/or a skin rash. If you experience a combination of these symptoms, contact your HCP immediately.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

Thrombotic thrombocytopenic purpura (TTP): LEMTRADA may cause blood clotting problems that can be fatal. Call your HCP right away if you experience symptoms such as: purplish spots on skin or in mouth due to bleeding under skin, yellowing of skin or whites of eyes (jaundice), feel tired or weak, very pale skin, fever, fast heart rate or short of breath, headache, speech changes, confusion, vision changes, seizure, low amount of urine or dark or bloody urine, stomach pain, nausea, vomiting, or diarrhea.

Autoimmune encephalitis (AIE): LEMTRADA may cause AIE, a brain disorder which may include symptoms that seem like an MS relapse. Call your HCP right away if you have any of the following symptoms: personality changes, mood changes, seeing things that are not there (hallucinations), agitation, short term memory loss, confusion, movement disorders, or seizures.

Bleeding disorder (acquired hemophilia A): LEMTRADA may cause acquired hemophilia A. Call your HCP right away if you have any of the following symptoms: bruising, nose bleeds, bleeding from a cut that may take longer than usual to stop, painful or swollen joints, blood in urine, dark or bloody stools.

Serious infections: LEMTRADA may cause you to have a serious infection while you receive and after receiving a course of treatment. Serious infections may include:

- **listeria.** People who receive LEMTRADA have an increased chance of getting a bacterial infection called listeria, which can lead to significant complications or death. Avoid foods that may be a source of listeria or make sure foods are heated well.
- **herpes viral infections.** Some people taking LEMTRADA have an increased chance of getting herpes viral infections. Take medicines as prescribed by your HCP to reduce your chances of getting these infections.
- **tuberculosis.** Your HCP should check you for tuberculosis before you receive LEMTRADA.
- **hepatitis.** People who are at high risk of, or are carriers of, hepatitis B (HBV) or hepatitis C (HCV) may be at risk of irreversible liver damage.

LEMTRADA[®]
alemtuzumab^{12mg} iv

IMPORTANT SAFETY INFORMATION (CONTINUED)

Serious infections (continued): These are not all the possible infections that could happen while on LEMTRADA. Call your HCP right away if you have symptoms of a serious infection such as fever or swollen glands. Talk to your HCP before you get vaccinations after receiving LEMTRADA. Certain vaccinations may increase your chances of getting infections.

Progressive multifocal leukoencephalopathy (PML): A rare brain infection that usually leads to death or severe disability has been reported with LEMTRADA. Symptoms of PML get worse over days to weeks. It is important that you call your doctor right away if you have any new or worsening medical problems that have lasted several days, including problems with thinking, eyesight, strength, balance, weakness on one side of your body, using your arms or legs.

Inflammation of the gallbladder without gallstones (acalculous cholecystitis): LEMTRADA may increase your chance of getting inflammation of the gallbladder without gallstones, a serious medical condition that can be life-threatening. Call your HCP right away if you have stomach pain or discomfort, fever, nausea, or are vomiting.

Swelling of lung tissue (pneumonitis): Some people have had swelling of the lung tissue while receiving LEMTRADA. Call your HCP right away if you have shortness of breath, cough, wheezing, chest pain or tightness, or are coughing up blood.

Before receiving LEMTRADA, tell your HCP if you:

- have bleeding, thyroid, or kidney problems
- have a recent history of infection
- are taking a medicine called Campath® (alemtuzumab)
- have received a live vaccine in the past 6 weeks before receiving LEMTRADA or plan to receive any live vaccines. Ask your HCP if you are not sure if your vaccine is a live vaccine.

- are pregnant or plan to become pregnant. LEMTRADA may harm your unborn baby. You should use birth control while receiving LEMTRADA and for 4 months after your course of treatment.
- are breastfeeding or plan to breastfeed. You and your HCP should decide if you should receive LEMTRADA or breastfeed.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LEMTRADA and other medicines may affect each other, causing side effects. Especially tell your HCP if you take medicines that increase your chance of getting infections, including medicines used to treat cancer or to control your immune system.

The most common side effects of LEMTRADA include:

- | | | |
|------------------------------------|-------------------------------|--|
| • rash | • upper respiratory infection | • diarrhea |
| • headache | • herpes viral infection | • sinus infection |
| • thyroid problems | • hives | • mouth pain or sore throat |
| • fever | • itching | • tingling sensation |
| • swelling of your nose and throat | • fungal infection | • dizziness |
| • nausea | • joint pain | • stomach pain |
| • urinary tract infection | • pain in your arms or legs | • sudden redness in face, neck, or chest |
| • feeling tired | • back pain | • vomiting |
| • trouble sleeping | | |

Tell your HCP if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of LEMTRADA.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

LEMTRADA[®]
alemtuzumab^{12mg} iv

FROM BEFORE TREATMENT TO MONITORING,
YOU'RE ON YOUR WAY

If you have any questions,
reach out to your *MS One to One*[®] Case Manager
or call **1-855-676-6326**

Please see Important Safety Information on pages 11-13 and full
[Prescribing Information/Medication Guide](#), including serious side effects.

sanofi

© 2022 Genzyme Corporation. All rights reserved.
Lemtrada, *MS One to One*, and Sanofi registered in U.S. Patent and Trademark Office.
MAT-US-2008398-v6.0-06/2022

LEMTRADA[®]
alemtuzumab^{12mg}
iv